arXiv:1210.2538v3 [hep-ph] 27 Oct 2012

KUNS-2420

Bare Higgs mass at Planck scale

Yuta Hamada,[*] Hikaru Kawai,[†] and Kin-ya Oda[‡]

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

October 30, 2012

Abstract

In a certain quantum gravity/string theory context, it is possible that not only the physical Higgs mass but also the bare one (and hence the radiative corrections as well) can vanish at the Planck/string scale. We compute one and two loop quadratic divergent contributions to the bare Higgs mass in terms of the bare couplings in the Standard Model (SM). We approximate the bare couplings, defined at the ultraviolet cutoff scale, by the $\overline{\text{MS}}$ ones at the same scale, which are evaluated by the two loop renormalization group equations for the Higgs mass around 126 GeV in the SM. We obtain the cutoff scale dependence of the bare Higgs mass, and examine where it becomes zero. We find that when we take the current central value for the top quark pole mass, 173 GeV, the bare Higgs mass vanishes if the cutoff is about 10^{23} GeV. With a $1.3\,\sigma$ smaller mass, 170 GeV, the scale can be of the order of the Planck scale.

---

[*] E-mail: `hamada@gauge.scphys.kyoto-u.ac.jp`

[†] E-mail: `hkawai@gauge.scphys.kyoto-u.ac.jp`

[‡] E-mail: `odakin@gauge.scphys.kyoto-u.ac.jp`

1 Introduction

The ATLAS and CMS experiments at the Large Hadron Collider (LHC) observed a particle at the 5σ confidence level (CL), which is consistent with the Standard Model (SM) Higgs boson with mass

$$m_H = \begin{cases} 126.0 \pm 0.4 \pm 0.4\,\text{GeV}, & \text{ATLAS [1]}, \\ 125.3 \pm 0.4 \pm 0.5\,\text{GeV}, & \text{CMS [2]}. \end{cases} \tag{1}$$

Such a relatively light Higgs boson is compatible with the electroweak precision data [3]. Furthermore, this value of Higgs mass allows the SM to be valid up to the Planck scale, within the unitarity, (meta)stability, and triviality bounds [4, 5, 6]. Up to now, there are no symptom of breakdown of the SM as an effective theory below the Planck scale.

On the other hand, if one wants to solve the Higgs mass fine tuning problem within a framework of quantum field theory, it would be natural to assume a new physics at around TeV scale. The supersymmetry is a possible solution to cancel the quadratic divergences in the Higgs mass, see e.g. Ref. [7]. However, a Higgs mass around 126 GeV requires some amount of fine tuning in the Higgs sector in the Minimal Supersymmetric Standard Model (MSSM), see e.g. Ref. [8]. Furthermore, no sign of supersymmetry has been observed at LHC so far [9].

Given the current experimental situation, it is important to examine a scenario in which the SM is valid towards a very high ultraviolet (UV) cutoff scale Λ. In such a case, the fine tuning of the Higgs mass would be achieved in a framework beyond the ordinary quantum field theory, as is necessary for the quartic and quadratic divergences of the cosmological constant, see e.g. Ref. [10].

In this paper, we compute the bare Higgs mass at Λ by taking into account one and two loop corrections in the SM. We will see that the bare mass can be zero if Λ is at around the Planck scale. We also discuss the possibility that a small non-vanishing bare mass in units of Λ can be canceled by threshold corrections in string theory.

This paper is organized as follows. In the next section, we explain our convention, and calculate the quadratic divergent contributions to the bare Higgs mass up to the two loop orders. In Section 3, we present a renormalization group equation (RGE) analysis in the SM and give our results for the Higgs quartic coupling at high scales. In Section 4, we examine how small the bare Higgs mass can be at the Planck scale, and show at what scale the bare Higgs mass vanishes. We vary α_s, m_H, and m_t^{pole} to see how the results are affected. The last section is the summary and discussions.

2 Bare Higgs mass

For the SM Higgs sector, we start from the bare Lagrangian of the following form in a fixed cutoff scheme with cutoff Λ:[1]

$$\mathcal{L} = (D_\mu \phi_B)^\dagger (D^\mu \phi_B) - m_B^2 \phi_B^\dagger \phi_B - \lambda_B (\phi_B^\dagger \phi_B)^2, \qquad \phi_B = \begin{pmatrix} \phi_B^+ \\ \phi_B^0 \end{pmatrix}. \tag{2}$$

The Planck scale is

$$M_{\text{Pl}} = \frac{1}{\sqrt{G_N}} = 1.22 \times 10^{19}\,\text{GeV}. \tag{3}$$

We take into account the SM couplings $g_Y, g_2, g_3, \lambda, y_t$ and neglect the others.

2.1 Bare mass in ϕ^4 theory

We compute the quadratic divergence in the bare Higgs mass. First we explain our procedure by taking the concrete evaluation for the ϕ^4 theory. From the bare Lagrangian

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \phi_B)^2 - \frac{m_B^2}{2}\phi_B^2 - \frac{\lambda_B}{4!}\phi_B^4, \tag{4}$$

we calculate the bare mass m_B^2 order by order in the loop expansion so that the physical mass is tuned to be zero[2]

$$m_B^2 = m^2_{B,\,\text{0-loop}} + m^2_{B,\,\text{1-loop}} + m^2_{B,\,\text{2-loop}} + \cdots . \tag{5}$$

At each order, we fix the bare mass as

$$m^2_{B,\,\text{0-loop}} = 0, \tag{6}$$

$m^2_{B,\,\text{1-loop}} + i \Bigg($



$\Bigg)\Bigg|_{k=0} = 0, \quad (7)$

$m^2_{B,\,\text{2-loop}} + i \Bigg($



$\Bigg)\Bigg|_{k=0} = 0. \quad (8)$

---

[1] We assume the form without higher dimensional operators for whole the bare SM Lagrangian, since these can be absorbed by the redefinition of the couplings when we are only interested in low energy observables.

[2] Precisely speaking, $m^2_{B,\,\text{0-loop}}$ corresponds to the physical mass times the wave function renormalization factor and is negligibly small compared to the UV cutoff scale.

The one loop integral in Eq. (7) is quadratically divergent and is proportional to

$$I_1 := \int \frac{d^4 p_E}{(2\pi)^4} \frac{1}{p_E^2}, \tag{9}$$

where p_E is a Euclidean four momentum. In the two loop computation (8), the momentum integrals in the third and fourth terms are, respectively,

$$J_2 := \int \frac{d^4 p_E}{(2\pi)^4} \frac{d^4 q_E}{(2\pi)^4} \frac{1}{p_E^4 q_E^2}, \tag{10}$$

$$I_2 := \int \frac{d^4 p_E}{(2\pi)^4} \frac{d^4 q_E}{(2\pi)^4} \frac{1}{p_E^2 q_E^2 (p_E + q_E)^2}. \tag{11}$$

The integral J_2 is infrared (IR) divergent: $J_2 \propto \Lambda^2 \ln(\Lambda/\mu_{\rm IR})$, but is canceled by the second term in Eq. (8) due to the lower order condition (7). Therefore we are left with only I_2 that does not suffer from the infrared divergence. This situation does not change in higher orders because a mass should not contain an IR divergence.

2.2 Bare mass in SM

Now let us fill the above prescription in the SM. In the following, we work in the symmetric phase $\langle \phi \rangle = 0$ as we are interested only in the quadratic divergent terms. In the evaluation of the Feynman diagrams, it is convenient to take the Landau gauge for all the $SU(3) \times SU(2) \times U(1)$ gauge fields. In this gauge, a diagram always vanishes if an external Higgs line is attached with a gauge boson propagator by a three point vertex:



$$= 0. \tag{12}$$

From the one loop diagrams we get the quadratic divergent integral I_1 again [15]

$$m^2_{B,\,\text{1-loop}} = -\left(6\lambda_B + \frac{3}{4} g^2_{YB} + \frac{9}{4} g^2_{2B} - 6 y^2_{tB}\right) I_1. \tag{13}$$

In Fig. 1, we present the two loop Feynman diagrams that do not vanish in the symmetric phase $\langle \phi \rangle = 0$ and in the Landau gauge. In the second row of Fig. 1, the last diagram cancels the divergences coming from the one-loop self energy of the internal Higgs propagators, as in Eq. (8).[3] All the momentum

---

[3] In practice, from each diagram containing a self energy correction, one subtracts a term that is obtained by setting the external momentum of its self energy to be zero. One may also apply this subtraction, though not necessary, for diagrams containing a vacuum polarization in order to simplify the calculation.



Figure 1: Non-vanishing two loop Feynman diagrams. Arrows are omitted. The dashed, solid, wavy, and dotted lines represent the scalar, fermion, gauge, and ghost propagators, respectively.

integrals can be recasted into either I_2 or J_2. We have explicitly checked that the coefficients of the infrared divergent integral J_2 cancel in each gauge invariant set of diagrams.

The two loop contribution to the bare Higgs mass at Λ becomes[4]

$$\begin{aligned} m^2_{B,\,\text{2-loop}} = -\Bigg\{ & 9y_{tB}^4 + y_{tB}^2\left(-\frac{7}{12}g_{YB}^2 + \frac{9}{4}g_{2B}^2 - 16g_{3B}^2\right) + \frac{77}{16}g_{YB}^4 + \frac{243}{16}g_{2B}^4 \\ & + \lambda_B\left(-18y_{tB}^2 + 3g_{YB}^2 + 9g_{2B}^2\right) - 10\lambda_B^2 \Bigg\} I_2. \end{aligned} \tag{14}$$

---

[4] As is mentioned in Ref. [15], while at the one-loop level only a restricted set of particles participates, on the two-loop level all kinds of particles up to the Planck mass enter in the discussion. We assume that there appear only SM degrees of freedom up to the UV cutoff scale.

This is one of our main results. Note that Eqs. (13) and (14) are minus the radiative corrections to the physical Higgs mass squared, see Eqs. (7) and (8).

In Section 4, we will examine whether the bare mass can vanish at a particular UV cutoff scale. For that purpose, we need to relate the integrals I_1 and I_2. This relation necessarily depends on the cutoff scheme. In particular, if the two loop contribution to the bare mass $m^2_{B,\,\text{2-loop}}$ becomes sizable compared to $m^2_{B,\,\text{1-loop}}$, the result suffers from a large theoretical uncertainty. We will verify that it is actually small. With this caution in mind, let us employ the following regularization:

$$\int d^4k_E \frac{1}{k_E^2} = \int_\varepsilon^\infty d\alpha \int d^4k_E\, e^{-\alpha k_E^2}, \tag{15}$$

which gives

$$I_1 = \frac{1}{\varepsilon}\frac{1}{16\pi^2}, \qquad\qquad I_2 = \frac{1}{\varepsilon}\frac{1}{(16\pi^2)^2}\ln\frac{2^6}{3^3} \simeq 0.005\, I_1. \tag{16}$$

When we employ a naive momentum cutoff by Λ, we get

$$I_1 = \frac{\Lambda^2}{16\pi^2}, \tag{17}$$

and hence we can regard $1/\varepsilon = \Lambda^2$.

2.3 Graviton effects

Let us estimate the graviton loop effects on the above obtained result. The graviton $h_{\mu\nu}$ in the metric

$$g_{\mu\nu} = \eta_{\mu\nu} + \frac{\sqrt{32\pi}}{M_{\text{Pl}}} h_{\mu\nu} \tag{18}$$

couples to the Higgs through the energy-momentum tensor:

$$\begin{aligned} T_{\mu\nu} &= \frac{2}{\sqrt{-g}}\frac{\delta}{\delta g^{\mu\nu}}\sqrt{-g}\mathcal{L} \\ &= (D_\mu\phi)^\dagger(D_\nu\phi) + (D_\nu\phi)^\dagger(D_\mu\phi) - g_{\mu\nu}\left[(D_\mu\phi)^\dagger(D^\mu\phi) - m_B^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2\right]. \end{aligned} \tag{19}$$

The most divergent contributions come from two derivative couplings. A one loop diagram containing such a graviton coupling vanishes because it necessarily picks up an external momentum, which is set to zero. Other contributions are at most logarithmically divergent. At the two loop level, diagrams involving internal graviton line that does not touch a Higgs external line give a form $\Lambda^4/M_{\text{Pl}}^2$. If the UV cutoff is much smaller than the Planck scale, this becomes negligible, and the higher loops become further insignificant. Indeed in perturbative string theory, higher loop corrections are proportional to powers of the string coupling constant g_s and become sub-leading. If the cutoff scale exceeds the Planck scale, we cannot neglect the graviton contributions.

3 SM RGE evolution toward Planck scale

In Section 4, we will approximate the bare coupling constants in the SM at the UV cutoff scale Λ by the running ones in the modified minimal subtraction ($\overline{\text{MS}}$) scheme at the same scale Λ; see Section 4.2 for its justification.

To get the $\overline{\text{MS}}$ running coupling constant, we apply the renormalization group equation (RGE) at the two loop order. For g_Y, g_2, g_3, and y_t, we use the ones in Ref. [16].[5] For the quartic coupling, we employ the one given in Ref. [17].[6] To be explicit,

$$
\begin{aligned}
\frac{dg_Y}{dt} &= \frac{1}{16\pi^2}\frac{41}{6}g_Y^3 + \frac{g_Y^3}{(16\pi^2)^2}\left(\frac{199}{18}g_Y^2 + \frac{9}{2}g_2^2 + \frac{44}{3}g_3^2 - \frac{17}{6}y_t^2\right), \\
\frac{dg_2}{dt} &= -\frac{1}{16\pi^2}\frac{19}{6}g_2^3 + \frac{g_2^3}{(16\pi^2)^2}\left(\frac{3}{2}g_Y^2 + \frac{35}{6}g_2^2 + 12g_3^2 - \frac{3}{2}y_t^2\right), \\
\frac{dg_3}{dt} &= -\frac{7}{16\pi^2}g_3^3 + \frac{g_3^3}{(16\pi^2)^2}\left(\frac{11}{6}g_Y^2 + \frac{9}{2}g_2^2 - 26g_3^2 - 2y_t^2\right), \\
\frac{dy_t}{dt} &= \frac{y_t}{16\pi^2}\left(\frac{9}{2}y_t^2 - \frac{17}{12}g_Y^2 - \frac{9}{4}g_2^2 - 8g_3^2\right) + \frac{y_t}{(16\pi^2)^2}\Bigg(-12y_t^2 + 6\lambda^2 - 12\lambda y_t^2 \\
&\quad + \frac{131}{16}g_Y^2y_t^2 + \frac{225}{16}g_2^2y_t^2 + 36g_3^2y_t^2 + \frac{1187}{216}g_Y^4 - \frac{23}{4}g_2^4 - 108g_3^4 - \frac{3}{4}g_Y^2g_2^2 + 9g_2^2g_3^2 + \frac{19}{9}g_3^2g_Y^2\Bigg), \\
\frac{d\lambda}{dt} &= \frac{1}{16\pi^2}\left(24\lambda^2 - 3g_Y^2\lambda - 9g_2^2\lambda + \frac{3}{8}g_Y^4 + \frac{3}{4}g_Y^2g_2^2 + \frac{9}{8}g_2^4 + 12\lambda y_t^2 - 6y_t^4\right) \\
&\quad + \frac{1}{(16\pi^2)^2}\Bigg\{ -312\lambda^3 + 36\lambda^2(g_Y^2 + 3g_2^2) - \lambda\left(\frac{629}{24}g_Y^4 - \frac{39}{4}g_Y^2g_2^2 + \frac{73}{8}g_2^4\right) \\
&\qquad + \frac{305}{16}g_2^6 - \frac{289}{48}g_Y^2g_2^4 - \frac{559}{48}g_Y^4g_2^2 - \frac{379}{48}g_Y^6 - 32g_3^2y_t^4 - \frac{8}{3}g_Y^2y_t^4 - \frac{9}{4}g_2^4y_t^2 \\
&\qquad + \lambda y_t^2\left(\frac{85}{6}g_Y^2 + \frac{45}{2}g_2^2 + 80g_3^2\right) + g_Y^2y_t^2\left(-\frac{19}{4}g_Y^2 + \frac{21}{2}g_2^2\right) \\
&\qquad - 144\lambda^2y_t^2 - 3\lambda y_t^4 + 30y_t^6\Bigg\},
\end{aligned}
\tag{20}
$$

where $t = \ln\mu$. Though we do not include the bottom and tau Yukawa couplings in this paper, we have checked that these are negligible within the precision that we work in.

We put the boundary condition for the RGE (20) according to Ref. [5]. The $\overline{\text{MS}}$ gauge coupling of $SU(3)$ is given by the three loop RGE running from m_Z

---

[5] We replace g_1 of the GUT normalization to $g_Y = \sqrt{3/5}\,g_1$ and rewrite the quartic coupling as $\lambda_{[16]} = 2\lambda$, where $\lambda_{[16]}$ is the one employed in Ref. [16].

[6] We use the arXiv version 2 of Ref. [17] with the replacements $g' = g_Y$, $g = g_2$, $h = y_t$, and $\lambda_{[17]} = 6\lambda$, where $\lambda_{[17]}$ is the quartic coupling employed in Ref. [17]. The RGE for λ in Ref. [16] becomes equal to that of Ref. [17], after correcting $-\frac{3}{2}g_2^4Y_4(S)$ to $-\frac{3}{2}g_2^4Y_2(S)$ and changing the part $\frac{229}{4} + \frac{50}{9}n_g$ to $\frac{229}{24} + \frac{50}{9}n_g$ in Eq. (A.17) in Ref. [16].

to $m_t^{\rm pole}$ and matching with six flavor theory as [5]

$$g_s(m_t^{\rm pole}) = 1.1645 + 0.0031\left(\frac{\alpha_s(m_Z) - 0.1184}{0.0007}\right) - 0.00046\left(\frac{m_t^{\rm pole}}{\rm GeV} - 173.15\right), \tag{21}$$

where $m_t^{\rm pole}$ is the pole mass of the top quark. The $\overline{\rm MS}$ quartic coupling at the top pole mass $m_t^{\rm pole}$ is given by taking into account the QCD and Yukawa two loop corrections [5]

$$\lambda(m_t^{\rm pole}) = 0.12577 + 0.00205\left(\frac{m_H}{\rm GeV} - 125\right) - 0.00004\left(\frac{m_t^{\rm pole}}{\rm GeV} - 173.15\right) \pm 0.00140_{\rm th}, \tag{22}$$

where m_H is the observed Higgs mass which we read off from Eq. (1) as

$$m_H = 125.7 \pm 0.6\,{\rm GeV}. \tag{23}$$

The $\overline{\rm MS}$ top Yukawa coupling at the scale $m_t^{\rm pole}$ is given by taking into account the QCD three loop, electroweak one loop, and $O(\alpha\alpha_s)$ two loop corrections [5]

$$\begin{aligned} y_t(m_t^{\rm pole}) = {} & 0.93587 + 0.00557\left(\frac{m_t^{\rm pole}}{\rm GeV} - 173.15\right) - 0.00003\left(\frac{m_H}{\rm GeV} - 125\right) \\ & - 0.00041\left(\frac{\alpha_s(m_Z) - 0.1184}{0.0007}\right) \pm 0.00200_{\rm th}. \end{aligned} \tag{24}$$

In a more recent work [6], it has been pointed out that the error in the top quark pole mass, consistently derived from the running one, is larger than that given in Ref. [5], 173.1 ± 0.7 GeV. The value obtained is [6]

$$m_t^{\rm pole} = 173.3 \pm 2.8\,{\rm GeV}, \tag{25}$$

which we will use in our analysis.

We plot the $\overline{\rm MS}$ running coupling constant $\lambda(\mu)$ in Fig. 2. As we increase the scale μ, the coupling λ first decreases due to the term $-6y_t^4$ and keeps to be small above $\mu = 10^{10}\,{\rm GeV}$ for a while. At further higher energies, y_t becomes smaller and λ starts to increase due to the contribution from $\frac{3}{8}g_Y^4$ which is not asymptotically free. At the intermediate scale, λ can become negative but it is shown that a metastability condition can be met even in this case [4, 5, 6]. The value of λ at the Planck scale $M_{\rm Pl}$ becomes consistent with Eq. (64) in Ref. [5]

$$\begin{aligned} \lambda(M_{\rm Pl}) = {} & -0.014 - 0.018\left(\frac{m_t^{\rm pole} - 173.3\,{\rm GeV}}{2.8\,{\rm GeV}}\right) + 0.002\left(\frac{\alpha_s(m_Z) - 0.1184}{0.0007}\right) \\ & + 0.002\left(\frac{m_H - 125.7\,{\rm GeV}}{0.6\,{\rm GeV}}\right) \pm 0.004_{\rm th}. \end{aligned} \tag{26}$$



Figure 2: Left: $\overline{\mathrm{MS}}$ running of the quartic coupling λ. The band corresponds to the 1σ deviation $m_t^{\mathrm{pole}} = 173.3 \pm 2.8\,\mathrm{GeV}$. Right: The scale $\mu_{\min}$ at which $\lambda(\mu)$ takes its minimum value, as a function of m_t^{pole}. In both panels, low energy inputs are given by the central values $\alpha_s(m_Z) = 0.1184$ and $m_H = 125.7\,\mathrm{GeV}$.

As we can see from the left panel in Fig. 2, the value of the quartic coupling stays around its minimum in $10^{15}\,\mathrm{GeV} \lesssim \mu \lesssim 10^{20}\,\mathrm{GeV}$. Therefore the minimum value of λ is also given by Eq. (26) within our precision. In the right panel in Fig. 2, we plot $\mu_{\min}$ at which the $\lambda(\mu)$ takes its minimum value. The central value $m_t^{\mathrm{pole}} = 173.3\,\mathrm{GeV}$ gives $\mu_{\min} = 4 \times 10^{17}\,\mathrm{GeV}$.

4 Bare Higgs mass at Planck scale

4.1 Numerical results

Now we can estimate the bare Higgs mass at the cutoff scale by substituting the $\overline{\mathrm{MS}}$ couplings derived in the previous section to the bare ones in the right hand sides of Eqs. (13) and (14).

In the left panel of Fig. 3, we plot the dependence of the bare Higgs mass-squared in units of $\Lambda^2/16\pi^2$ on the UV cutoff scale Λ:

$$\frac{m_B^2}{\Lambda^2/16\pi^2} = \frac{m^2_{B,\,\text{1-loop}}}{I_1} + \frac{m^2_{B,\,\text{2-loop}}}{I_2}\frac{I_2}{I_1}, \tag{27}$$

where we have taken $I_2/I_1 = 0.005$ as in Eq. (16). In the figure, we can see that the bare mass m_B^2 monotonically decreases when one increase Λ.

We obtain the UV cutoff scale at which the bare mass m_B^2 becomes zero:

$$\log_{10}\frac{\Lambda|_{m_B^2=0}}{\mathrm{GeV}} = 23.5 + 3.3\left(\frac{m_t^{\mathrm{pole}} - 173.3\,\mathrm{GeV}}{2.8\,\mathrm{GeV}}\right) - 0.2\left(\frac{m_H - 125.7\,\mathrm{GeV}}{0.6\,\mathrm{GeV}}\right)$$
$$- 0.4\left(\frac{\alpha_s(m_Z) - 0.1184}{0.0007}\right) \pm 0.4_{\mathrm{th}}. \tag{28}$$



Figure 3: Left: The bare Higgs mass m_B^2 in units of $\Lambda^2/16\pi^2$ vs the UV cutoff scale Λ. The blue (narrower) and pink (wider) bands represent the one and two sigma deviations of m_t^{pole}, respectively. Right: The UV cutoff scale at which the bare mass m_B^2 becomes zero as a function of m_t^{pole}. The solid (dashed) line corresponds to the scale where m_B^2 ($m_{B,\,\text{1-loop}}^2$) becomes zero. In both panels, we have taken the central values $\alpha_s(m_Z) = 0.1184$ and $m_H = 125.7\,\mathrm{GeV}$.

In the right panel of Fig. 3, we plot this quantity as a function of the top quark pole mass for the central values of $\alpha_s(m_Z)$ and m_H, without referring to the linear approximation (28).

We show an approximate formula for the bare Higgs mass when the cutoff is at the Planck scale, $\Lambda = M_{\mathrm{Pl}}$:

$$m_B^2 = \left[0.22 + 0.18\left(\frac{m_t^{\mathrm{pole}} - 173.3\,\mathrm{GeV}}{2.8\,\mathrm{GeV}}\right) - 0.02\left(\frac{\alpha_s(m_Z) - 0.1184}{0.0007}\right) - 0.01\left(\frac{m_H - 125.7\,\mathrm{GeV}}{0.6\,\mathrm{GeV}}\right) \pm 0.02_{\mathrm{th}}\right]\frac{M_{\mathrm{Pl}}^2}{16\pi^2}. \tag{29}$$

This is one of our main results. We verify that the two loop correction (14) can be safely neglected: $m_{B,\,\text{2-loop}}^2 \simeq -0.005\,M_{\mathrm{Pl}}^2/16\pi^2$ within the scheme (16), as advertised before. In Fig. 4, we plot the bare Higgs mass-squared in units of $M_{\mathrm{Pl}}^2/16\pi^2$ as a function of m_t^{pole} for the central values of $\alpha_s(m_Z)$ and m_H, without referring to the linear approximation (29). For comparison, we also plot the quartic coupling λ at the Planck scale.

From Fig. 4 we see that the bare Higgs mass becomes zero if $m_t^{\mathrm{pole}} = 169.8\,\mathrm{GeV}$, while the quartic coupling $\lambda(M_{\mathrm{Pl}})$ vanishes if $m_t^{\mathrm{pole}} = 171.2\,\mathrm{GeV}$, when we take the central values for $\alpha_s(m_Z)$ and m_H. See Refs. [11, 12, 13] for arguments supporting the vanishing quartic coupling at a cutoff scale, see also Ref. [14]. There is no low energy parameter set within two sigma that makes both the quartic coupling and the bare mass vanish simultaneously at the Planck scale. This might suggest an existence of a small threshold effect from an underlying UV complete theory.



Figure 4: The blue solid (dashed) line corresponds to the 1 plus 2 loop (1-loop) bare mass m_B^2 ($m_{B,\,\text{1-loop}}^2$) in units of $M_{\rm Pl}^2/16\pi^2$ for $\Lambda = M_{\rm Pl}$. For comparison, we also plot the quartic coupling λ at the Planck scale by the red dotted line. The central values $\alpha_s(m_Z) = 0.1184$ and $m_H = 125.7\,\text{GeV}$ are used.

4.2 Cutoff vs $\overline{\text{MS}}$

We have approximated the bare coupling constants in the SM by the running ones in the $\overline{\text{MS}}$ scheme at Λ. The resulting error can be evaluated once the cutoff scheme is explicitly specified.

More concretely, let us first express the $\overline{\text{MS}}$ couplings at a scale μ in terms of the bare couplings defined at the cutoff scale Λ:

$$\begin{aligned}\lambda^i_{\overline{\text{MS}}}(\mu) &= \lambda^i_B + \sum_{jk} c^{ijk}(\mu/\Lambda)\ \lambda^j_B \lambda^k_B + O(\lambda_B^3),\\ c^{ijk}(x) &:= f^{ijk} + b^{ijk}\ln x + O(x^2),\end{aligned} \tag{30}$$

where b^{ijk} is the coefficient in the one loop beta function and f^{ijk} is the finite part from the one loop diagrams. $\{\lambda^i_{\overline{\text{MS}}}\}_{i=1,\dots,5}$ ($\{\lambda^i_B\}_{i=1,\dots,5}$) stands for the $\overline{\text{MS}}$ (bare) couplings of the SM: $\{g_Y^2, g_2^2, g_3^2, y_t^2, \lambda\}$ ($\{g_{YB}^2, g_{2B}^2, g_{3B}^2, y_{tB}^2, \lambda_B\}$). In our case, the two loop corrections in the RGE at high scales are small compared to the one loop order, which indicates that the two loop terms $O(\lambda_B^3)$ in Eq. (30) are negligible as we can take μ that satisfies both

$$\mu \ll \Lambda, \qquad \left|\frac{\lambda^i_{\overline{\text{MS}}}}{16\pi^2}\ln(\mu/\Lambda)\right| \ll 1, \tag{31}$$

simultaneously. Thus we have

$$\lambda^i_{\overline{\text{MS}}}(\mu) = \lambda^i_B + \sum_{jk}\left(f^{ijk} + b^{ijk}\ln\frac{\mu}{\Lambda}\right)\lambda^j_B\lambda^k_B. \tag{32}$$

On the other hand, from the RGE, we get

$$\lambda^i_{\overline{\rm MS}}(\Lambda) = \lambda^i_{\overline{\rm MS}}(\mu) + \sum_{jk} b^{ijk} \lambda^j_{\overline{\rm MS}}(\mu)\, \lambda^k_{\overline{\rm MS}}(\mu) \ln \frac{\Lambda}{\mu}. \tag{33}$$

From Eqs. (32) and (33), we obtain

$$\lambda^i_{\overline{\rm MS}}(\Lambda) = \lambda^i_B + \sum_{jk} f^{ijk} \lambda^j_B \lambda^k_B, \tag{34}$$

which gives the relation between the bare and the $\overline{\rm MS}$ couplings at the same scale.

With the above correction, the formula for the bare Higgs mass is modified by

$$\Delta m_B^2 = -\sum_{ijk} a^i f^{ijk} \lambda^j_{\overline{\rm MS}}(\Lambda)\, \lambda^k_{\overline{\rm MS}}(\Lambda), \tag{35}$$

where a^i are the coefficients in the one loop bare Higgs mass $m_B^2 = \sum_i a^i \lambda^i_B$ in Eq. (13), and are proportional to I_1. The scale at which the bare Higgs mass vanishes $\Lambda|_{m_B^2=0}$ is changed to $\Lambda|_{m_B^2=0}\, e^{\delta t}$, where

$$\delta t = \frac{\sum_{ijk} a^i f^{ijk} \lambda^j_{\overline{\rm MS}}(\Lambda)\, \lambda^k_{\overline{\rm MS}}(\Lambda)}{\sum_{ijk} a^i b^{ijk} \lambda^j_{\overline{\rm MS}}(\Lambda)\, \lambda^k_{\overline{\rm MS}}(\Lambda)}. \tag{36}$$

Generically f^{ijk} are of the same order as b^{ijk} and hence the correction due to the replacement of the bare couplings by the $\overline{\rm MS}$ ones, Δm_B^2, is as small as the two loop corrections. Since δt is of order unity, the ambiguity for the scale $\Lambda|_{m_B^2=0}$ would be at most $e^{\delta t} \lesssim 10$.

5 Summary and discussions

We have presented a scheme where the quadratic divergence of the bare Higgs mass is computed in terms of the bare couplings at a UV cutoff scale Λ. Using it, we have obtained the bare Higgs mass up to the two loop order in the SM. This calculation has been made easier by working in the symmetric phase $\langle\phi\rangle = 0$ and in the Landau gauge. We have checked that all the IR divergent terms, which are proportional to $\Lambda^2 \ln(\Lambda/\mu_{\rm IR})$, cancel out as is expected. Approximating the bare couplings at Λ by the corresponding $\overline{\rm MS}$ ones at the same scale, we can examine whether the quadratic divergence in the bare Higgs mass vanish or not. To get the $\overline{\rm MS}$ couplings at high scales, we employ the two loop RGE in the SM. We have found that it is indeed the case if the top quark mass is

$m_t^{\text{pole}} = 169.8\,\text{GeV}$, which is $1.3\,\sigma$ smaller than the current central value.[7] One might find it intriguing that this value is close to $m_t^{\text{pole}} = 171.2\,\text{GeV}$ that gives vanishing quartic coupling at M_{Pl}.

There are several arguments [10, 11] suggesting that all the couplings, including the bare Higgs mass m_B^2, should vanish at a certain UV cutoff scale. It is a curious fact that the scale of the vanishing bare Higgs mass m_B^2 and that for the quartic coupling λ are quite close to each other and to the Planck scale. This result, obtained solely by using the SM, may indicate that the SM is valid up to the Planck scale and is a direct consequence of an underlying physics there.

If we take all the central values for m_t^{pole}, $\alpha_s(m_Z)$, and m_H, then the cancellation occurs not at the Planck scale but at a scale around $\Lambda \sim 10^{23}\,\text{GeV}$. This may hint a new physics around that scale. In this case however, we need to take the graviton effects into account, as is discussed in Section 2.3.

There can be a different interpretation for the small bare Higgs mass m_B^2 left at the Planck scale. It might appear as a threshold correction in string theory. In string theory, the tree level masses of the particles are quantized by $m_s := (\alpha')^{-1/2}$, and therefore the Higgs mass is zero at the tree level. The threshold effect from integrating out the massive stringy excitations is obtained by computing insertions of two Higgs emission vertices with zero external momenta into the world sheet. The result would become

$$m_B^2 \sim C \frac{g_s^2}{16\pi^2} m_s^2, \tag{37}$$

where C is a model dependent constant. This calculation can be performed for a concrete model such as the orbifold and fermionic constructions in heterotic string. This work will be presented in a separate publication.

We comment on the case where the UV completion of the SM appears as a supersymmetry. When the supersymmetry is softly broken, there cannot be any quadratic divergence and our study does not apply. In the case of the split/high-scale supersymmetry [18, 12] it is possible to perform a parallel analysis to the current one, which will be shown elsewhere.

If we assume the see-saw mechanism, the right-handed neutrinos are introduced above an intermediate scale M_R. Our analysis corresponds to the case where M_R is small enough that all the neutrino Dirac Yukawa couplings are negligible $y_D \lesssim 10^{-2}$. This condition implies $M_R \lesssim 10^{10}\,\text{GeV}$ for the neutrino mass $m_\nu \sim y_D^2 v^2/M_R \sim 0.1\,\text{eV}$. It would be interesting to extend our analysis to include larger Dirac Yukawa couplings for $M_R \gtrsim 10^{10}\,\text{GeV}$.

---

[7] The vanishing of the quadratic divergence does not immediately indicate that the bare Higgs mass is exactly zero. Our result does not exclude logarithmically divergent corrections such as $m_H^2 \ln(\Lambda/m_H)$ as well as finite ones. If the quadratic divergence indeed vanishes exactly for some reason, then such corrections become important. It would be interesting to study them.

Acknowledgement

We greatly appreciate valuable discussions with Satoshi Iso and useful information from him, without which this study would have never been started. This work is in part supported by the Grant-in-Aid for Scientific Research Nos. 22540277 (HK), 23104009, 20244028, and 23740192 (KO) and for the Global COE program "The Next Generation of Physics, Spun from Universality and Emergence."

References

[1] G. Aad *et al.* [ATLAS Collaboration], *"Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,"* Phys. Lett. B **716** (2012) 1 [arXiv:1207.7214 [hep-ex]].

[2] S. Chatrchyan *et al.* [CMS Collaboration], *"Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC,"* Phys. Lett. B **716** (2012) 30 [arXiv:1207.7235 [hep-ex]].

[3] J. Beringer *et al.* [Particle Data Group Collaboration], *"Review of Particle Physics (RPP),"* Phys. Rev. D **86** (2012) 010001.

[4] M. Holthausen, K. S. Lim and M. Lindner, *"Planck scale Boundary Conditions and the Higgs Mass,"* JHEP **1202** (2012) 037 [arXiv:1112.2415 [hep-ph]];
F. Bezrukov, M. Y. Kalmykov, B. A. Kniehl and M. Shaposhnikov, *"Higgs boson mass and new physics,"* arXiv:1205.2893 [hep-ph];
I. Masina, *"The Higgs boson and Top quark masses as tests of Electroweak Vacuum Stability,"* arXiv:1209.0393 [hep-ph].

[5] G. Degrassi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori and A. Strumia, *"Higgs mass and vacuum stability in the Standard Model at NNLO,"* JHEP **1208** (2012) 098 [arXiv:1205.6497 [hep-ph]].

[6] S. Alekhin, A. Djouadi and S. Moch, *"The top quark and Higgs boson masses and the stability of the electroweak vacuum,"* Phys. Lett. B **716** (2012) 214 [arXiv:1207.0980 [hep-ph]].

[7] S. P. Martin, *"A Supersymmetry primer,"* In *Kane, G.L. (ed.): Perspectives on supersymmetry II* 1-153 [hep-ph/9709356].

[8] H. Baer, V. Barger, P. Huang and X. Tata, *"Natural Supersymmetry: LHC, dark matter and ILC searches,"* JHEP **1205** (2012) 109 [arXiv:1203.5539 [hep-ph]].

[9] The ATLAS Collaboration, *"Search for new phenomena using large jet multiplicities and missing transverse momentum with ATLAS in 5.8 fb^{-1} of $\sqrt{s} = 8$ TeV proton-proton collisions,"* ATLAS-CONF-2012-103;

"Search for supersymmetry at $\sqrt{s} = 8$ TeV in final states with jets, missing transverse momentum and one isolated lepton," ATLAS-CONF-2012-104; *"Search for Supersymmetry in final states with two same-sign leptons, jets and missing transverse momentum with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV,"* ATLAS-CONF-2012-105; *"Search for squarks and gluinos with the ATLAS detector using final states with jets and missing transverse momentum at $\sqrt{s} = 8$ TeV,"* ATLAS-CONF-2012-109; The CMS Collaboration, *"Search for supersymmetery in final states with missing transverse momentum and 0, 1, 2, or ≥3 b jets in 8 TeV pp collisions"* CMS-PAS-SUS-12-016; *"Search for New Physics in Events with a Z Boson and Missing Transverse Energy,"* CMS-PAS-SUS-12-017; *"Search for supersymmetry in events with photons and missing energy,"* CMS-PAS-SUS-12-018.

[10] A. D. Linde, *"The Universe Multiplication And The Cosmological Constant Problem,"* Phys. Lett. B **200** (1988) 272; S. R. Coleman, *"Why There Is Nothing Rather Than Something: A Theory of the Cosmological Constant,"* Nucl. Phys. B **310** (1988) 643; S. Weinberg, *"The Cosmological Constant Problem,"* Rev. Mod. Phys. **61** (1989) 1.

[11] C. D. Froggatt and H. B. Nielsen, *"Standard model criticality prediction: Top mass 173 ± 5-GeV and Higgs mass 135 +- 9-GeV,"* Phys. Lett. B **368** (1996) 96 [hep-ph/9511371]; B. Stech in *"Proceedings to the Workshop at Bled, Slovenia, 29 June - 9 July 1998: What comes beyond the Standard Model,"* Ljubljana, Slovenia: DMFA (1999) 73 p [hep-ph/9905357]; K. A. Meissner and H. Nicolai, *"Conformal Symmetry and the Standard Model,"* Phys. Lett. B **648** (2007) 312 [hep-th/0612165]; *"Effective action, conformal anomaly and the issue of quadratic divergences,"* Phys. Lett. B **660** (2008) 260 [arXiv:0710.2840 [hep-th]]; S. Iso, N. Okada and Y. Orikasa, *"Classically conformal $B-L$ extended Standard Model,"* Phys. Lett. B **676** (2009) 81 [arXiv:0902.4050 [hep-ph]]; *"The minimal $B-L$ model naturally realized at TeV scale,"* Phys. Rev. D **80** (2009) 115007 [arXiv:0909.0128 [hep-ph]]; H. Kawai and T. Okada, *"Solving the Naturalness Problem by Baby Universes in the Lorentzian Multiverse,"* Prog. Theor. Phys. **127** (2012) 689 [arXiv:1110.2303 [hep-th]].

[12] L. J. Hall and Y. Nomura, *"A Finely-Predicted Higgs Boson Mass from A Finely-Tuned Weak Scale,"* JHEP **1003** (2010) 076 [arXiv:0910.2235 [hep-ph]].

[13] M. Shaposhnikov and C. Wetterich, *"Asymptotic safety of gravity and the Higgs boson mass,"* Phys. Lett. B **683** (2010) 196 [arXiv:0912.0208 [hep-th]].

[14] R. Foot, A. Kobakhidze and R. R. Volkas, *"Electroweak Higgs as a pseudo-Goldstone boson of broken scale invariance,"* Phys. Lett. B **655**, 156 (2007) [arXiv:0704.1165 [hep-ph]];
R. Foot, A. Kobakhidze, K. L. McDonald and R. R. Volkas, *"A Solution to the hierarchy problem from an almost decoupled hidden sector within a classically scale invariant theory,"* Phys. Rev. D **77**, 035006 (2008) [arXiv:0709.2750 [hep-ph]].

[15] M. J. G. Veltman, *"The Infrared - Ultraviolet Connection,"* Acta Phys. Polon. B **12** (1981) 437.

[16] H. Arason, D. J. Castano, B. Keszthelyi, S. Mikaelian, E. J. Piard, P. Ramond and B. D. Wright, *"Renormalization group study of the standard model and its extensions. 1. The Standard model,"* Phys. Rev. D **46** (1992) 3945.

[17] C. Ford, I. Jack and D. R. T. Jones, *"The Standard model effective potential at two loops,"* Nucl. Phys. B **387** (1992) 373 [Erratum-ibid. B **504** (1997) 551] [hep-ph/0111190].

[18] N. Arkani-Hamed and S. Dimopoulos, *"Supersymmetric unification without low energy supersymmetry and signatures for fine-tuning at the LHC,"* JHEP **0506** (2005) 073 [hep-th/0405159].